

INTRA-OFFICE MEMORANDUM

25X1X6

TO:

DATE: April 1, 1953

FROM:

SUBJECT: New Products - March, 1953

Domestic

Bicillin, All-purpose Injection

Description: An injectable penicillin combining in a single preparation, Bicillin, Procaine Penicillin and Potassium Penicillin.

Indications: Used for treatment against most penicillin-susceptible organisms. It "provides new penicillin blood levels with high initial peak, prolonged maintenance plateau and extended therapeutic and prophylactic levels."

25X1X6

Form Supplied: 1-dose vials.

Source:

Cortone with Bacitracin

Description: Ophthalmic ointment in tubes containing 15 mg. of "Cortone" (cortisone acetate, Merck) and 1,000 units of bacitracin per gram in a petrolatum base.

25X1X6

Indications: For treatment in certain diseases of the eye.

Form Supplied: Tubes of 2.5 grams.

Source:

Du-Biotic

Description: Intranasal
Contains phenylephrine hydrochloride, neomycin and bacitracin.

Indications: Subacute and chronic sinusitis; bacterial nasal infections.

Form Supplied: Dropper bottles containing the buffered active ingredients as a dry powder. Constituted by adding 15cc. of distilled water.

Troches

Description: Contains propesin, neomycin and bacitracin

Indications: Tonsillitis, pharyngitis, superficial mouth infections.

Form Supplied: In vials of 15 troches

Source: 25X1X6

Mycitracin

Description: A bland ointment containing neomycin and bacitracin.

Indications: Treatment of impetigo, barber's itch, infectious eczematoid dermatitis, infected ulcers, boils and other superficial pyogenic infections of the skin.

Administration: Topically, two to five times a day, with or without bandages.

Form Supplied: One-half ounce tubes.

Source: The Upjohn Co., Kalamazoo, Michigan

-2-

Neolin (Also see Sulfa-Neolin)

Description: Suspension containing di-benzylethylenediamine
dipenicillin G.
Indications: Penicillin-susceptible diseases including infections
due to pneumococcus, hemolytic streptococcus,
gonococcus and staphylococcus.
Administration: 1 teaspoonful four times daily in average adult
25X1X6 dose.
Form Supplied: Packages of 60 cc.
Source: [REDACTED]

Sulfa-Neolin

Description: A suspension containing sulfadiazine, sulfamerazine
and sulfamethazine and di-benzylethylenediamine
dipenicillin G.
For information as to administration, indications
etc. see Neolin.

-3-

Foreign

Adiazine a la Penicilline

Description: Each tablet contains:
 Sulfadiazine 0.50 g.
 Potassium Penicillin G 100,000 u.
 Calcium Carbonate 0.25 g.

Indications: For treatment of any infection sensitive to penicillin or sulfa. (A complement of Penicillinotherapie Parenterale).

25X1X6

Form Supplied: Bottles of 4 and 12 tablets

Source: [REDACTED] A.

Borocillina

Description: Each tablet contains:
 Penicillin 5,000 U.I.

Indications: Used in treatment of infections of the mouth and throat.

25X1X6

Administration: Dissolve in mouth slowly. 1 tablet every 1 to 2 hours.

Source: [REDACTED]

Entera-strept

Description: A tablet containing streptomycin.

Indications: Children's dyspepsia, sulfonamide-resistant bacteria and as a prophylaxis before operations.

Source: 25X1X6 [REDACTED]

Especilin - methiscol

Description: Each vial contains:
 Penicillin G. Procaine 300,000 u.ox.
 Penicillin G. Sodium 100,000 u.ox.
 Dihydrostreptomycin 0.5 g.
 Each accompanying vial of anti-toxic diluent contains:
 Acetyl Methiomine 0.780 g.
 Choline hydrochloride 0.300 g.
 Inositol 0.060 g.
 Water - double distilled 3 cm³

Indications: Used in surgery either pre- or post-operatively.

Source: Manufacturer not given

Distributors: 25X1X6 [REDACTED]

Formocilina

Description: Sulfamerazine
 Penicillin
 Streptomycin 25X1X6
 Chlorophyll

Indications: For treatment of infections in the oral cavity.

Source: [REDACTED]

-4-

Neosulfan

Description:

Contains tyrothrycin and sulfa drugs. Used to prevent and cure nose and throat infections.

25X1X6 [REDACTED]

Nocorin

Description:

Each lozenge contains:

Tyrothrycin

.001 G.

Benzocaine

.003 G.

25X1X6

Vitamin C (equal to 1,000 U.I. of ascorbic acid)

Indications:

For infections of the mouth and throat.

Source:

Omnamycin

Description:

Each vial contains:

Penicillin G

100,000 U.

Procaine Penicillin G

400,000 U.

Dihydrostreptomycin Sulfate

.5 Gm.

25X1X6

Omnadin (dry preparation) with diluent
(See December summary for description)

Source:

Paratebin

Description:

Each vial contains:

Oxyprocaine Penicillin

400,000 U.

Dihydrostreptomycin

1 Gm.

Indications:

For intensification of tuberculosis therapy.

Administration:

Intramuscular

Form Supplied:

Boxes of 1 vial with 1 ampule of 2 cc.

distilled water.

Source: 25X1X6 [REDACTED]

Pendex

Description:

A stable aqueous solution containing 1,500 units of penicillin per ml. with "paredrinex".

Indications:

25X1X6

Used in the intranasal treatment of sinusitis, rhinitis and nasopharyngitis.

Form Supplied:

15 ml. bottles with dropper.

Source:

Peneucin

Description:

Each gram contains 1,000 units of Penicillin

Indications:

Used in treatment of staphylococcus and streptococcus infections of the skin i.e. sycosis barbae, impetigo, secondarily infected eczema.

Form Supplied:

Tubes of 30 grams

Source:

25X1X6 [REDACTED]

-5-

Tirotrisol

Ginecologico

A vaginal antiseptic containing tyrothrycin, orthoxy-quinoline and lactic acid. It has an antibiotic and ph lowering action.

Nasal Drops

Used as antiseptic and to relieve congestion in the nose and throat. Contains tyrothrycin and desoxyephedrine, in an aqueous, isotonic, glucose solution of physiological pH. No oil or menthol.

Tablets

Contains tyrothrycin in an excipient which permits the antibiotic action to be maintained in the mouth and throat for 15 to 20 minutes.

Solution

For antibiotic action in mouth and throat. Contains tyrothrycin in a water-glycerin-alcohol solution with sodium salicylate and borate.

Ointment

Greaseless atoxic of physiological pH. Contains tyrothrycin, ethyl-urethane and bismuth sulfathiazole. An antiseptic used in the cicatrization and regeneration of tissues.

Source:

Trachicillin

25X1X6 Penicillin tablets used in treating angina, tonsillitis, gingivitis, stomatitis and after tonsillectomy and tooth extractions.

Source:

Trilcin Suspension

Description:

Each 4 ml. provides:

Sulfamethazine

0.12 gm.

Sulfadiazine

0.12 gm.

Sulfamerazine

0.09 gm.

Penicillin G. Potassium

100,000 I.U.

(Issued as the dry salt to be added

to the suspension at the time of dispensing)

Indications:

Infections caused by organisms which are sensitive to penicillin and sulphonamides.

Form Supplied:

Bottles of 48 ml.

Source:

25X1X6

Isobiocin

Maggioni & Co., Ltd. of Milan, Italy is marketing an injectable form of isonicotinic acid. Isobiocin is supplied in boxes of five 2 cc. ampules (100 mg.) as well as in bottles of 50 mg. tablets for oral therapy.